

## SUPPLEMENTARY INFORMATION

### Immunological response to nitroglycerin-loaded shear-responsive liposomes *in vitro* and *in vivo*

Marzia Buscema<sup>a</sup>, Sofiya Matviyiv<sup>a</sup>, Tamás Mészáros<sup>b,c</sup>, Gabriela Gerganova<sup>a</sup>,  
Andreas Weinberger<sup>d</sup>, Ute Mettal<sup>d</sup>, Dennis Mueller<sup>d</sup>, Frederik Neuhaus<sup>d</sup>, Etienne Stalder<sup>d</sup>,  
Takashi Ishikawa<sup>e</sup>, Rudolf Urbanics<sup>c</sup>, Till Saxer<sup>f</sup>, Thomas Pfohl<sup>a</sup>, János Szebeni<sup>b,c</sup>,  
Andreas Zumbuehl<sup>d</sup> and Bert Müller<sup>a</sup>

<sup>a</sup>Biomaterials Science Center, Department of Biomedical Engineering, University of Basel, Allschwil, Switzerland

<sup>b</sup>Nanomedicine Research and Education Center, Semmelweis University Budapest, Hungary

<sup>c</sup>SeroScience Ltd, Budapest, Hungary


<sup>d</sup>Department of Chemistry, University of Fribourg, Fribourg, Switzerland

<sup>e</sup>Paul Scherrer Institute (PSI), Villigen, Switzerland

<sup>f</sup>Cardiology Division, University Hospital of Geneva, Geneva, Switzerland

\* Corresponding author at: Biomaterials Science Center, Department of Biomedical Engineering, University of Basel, 4123 Allschwil, Gewerbestrasse 14, Switzerland, Tel: +41 61 207 54 30

E-mail address: bert.mueller@unibas.ch (B. Müller).




**Fig. S1. Time point *in vitro* experiment.** SC5b-9 concentration of six human sera incubated at a temperature of 37 °C with the positive, negative controls and liposomes at high lipid content (A) and at low lipid content (B). The data are shown as the mean value among the six donors. The reaction was terminated after 5, 10, 20, and 40 minutes. The data were sorted as A and B, in order to visualize clearly the SC5b-9 level of the Pad-PC-Pad-based liposomes at high and low lipid content.


**Table S2**

**Temporal sequences of liposomal injections in the six pigs, from left to right.** L1' was administrated in three liposomal concentrations, while L2' and L3' in a single concentration. For each injection, the pulmonary arterial pressure, PAP, and systemic arterial pressure, SAP, and heart rate, HR, were recorded and displayed in Figures S3 to S8.


Pig ID	Injection							
	#1	#2	#3	#4	#5	#6	#7	#8
1	1/3 × L1'	L1'	3 × L1'	L2'	L3'	Perlinganit	zymosan	
2	saline	3 × L1'	L2'	L3'	Perlinganit	1/3 × L1'	zymosan	
3	saline	3 × L1'	L1'	L2'	L3'	Perlinganit	zymosan	
4	saline	L1'	3 × L1'	L2'	L3'	Perlinganit	zymosan	
5	saline	L1'	3 × L1'	L3'	Perlinganit	zymosan		
6	saline	3 × L1'	L1'	1/3 × L1'	L2'	L3'	Perlinganit	zymosan




**Fig. S3.** Fig 1 (21 kg male)—Summary: Monitoring of PAP, SAP, and HR changes after injection; dashed line denotes injection time.




**Fig. S4.** Fig 2 (20 kg male)—Summary: Monitoring of PAP, SAP, and HR changes after injection; dashed line denotes injection time.




**Fig. S5.** Fig 3 (22 kg male)—Summary: Monitoring of PAP, SAP, and HR changes after injection; dashed line denotes injection time.



**Fig. S6.** Fig 4 (19 kg male)—Summary: Monitoring of PAP, SAP, and HR changes after injection; dashed line denotes injection time.



**Fig. S7.** Pig 5 (19 kg male)—Summary: Monitoring of PAP, SAP, and HR changes after injection; dashed line denotes injection time.



**Fig. S8.** Fig 6 (24 kg male)—Summary: Monitoring of PAP, SAP, and HR changes after injection; dashed line denotes injection time.